

# GeneProof Legionella pneumophila PCR Kit



### In vitro diagnostic medical device

The kit has been manufactured according to EC Directive 98/79/EC as an *in vitro* diagnostic medical device and it has been designed for professional use in specialized clinical and research laboratories.

### KIT CONTENT

	IS inc	ISIN Version cluded in the Mas	terMix	-	ISEX Version oplied in a separat plation and PCR ir	
REF	LP/ISIN/025	LP/ISIN/050	LP/ISIN/100	LP/ISEX/025	LP/ISEX/050	LP/ISEX/100
KLI	25 rxn	50 rxn	100 rxn	25 rxn	50 rxn	100 rxn
MasterMix Legionella pneumophila	1x750 μl	2x750 μl	4x750 μl	1x750 μl	2x750 μl	4x750 μl
<b>Positive Control</b> Legionella pneumophila	1x200 µl	1x200 μl	2x200 μl	1x200 μl	1x200 μl	2x200 μl
<b>Internal Standard</b> Legionella pneumophila	-	-	-	1x1000 μl	1x1000 µl	2x1000 μl

## STORAGE AND TRANSPORTATION CONDITIONS

The kits should be transported and stored at temperatures between -85 °C and -10 °C. The kit will remain stable at least until the expiry date printed on the package, if the storage temperature is kept. Repeated freezing and thawing of the kit components may result in lower detection quality.

### TECHNICAL SPECIFICATION

Target sequence	DNA conservative region encoding 16S rRNA		
Specificity	L. pneumophila (strain Corby, strain Lens, strain Paris, subsp. Pneumophila, subsp. Fraseri, subsp. Pascullei)		
Sensitivity (LOD)	reaches 1.398 copies/ $\mu$ l with the probability of 95 %		
Sample types	bronchoalveolar lavage, sputum		
Quality Control	regularly tested by QCMD and Instand e.V. External Quality Assessment Panels		



### METHOD PRINCIPLES

The PCR kit is designed for Legionella pneumophila detection by the Polymerase Chain Reaction (PCR) method. The L. pneumophila detection is based on the amplification of a sequence 16S rDNA gene specific for L. pneumophila and measuring the amplification product concentration using the PCR process and fluorescence labelled probes. L. pneumophila presence is indicated by FAM fluorophore fluorescence growth. An Internal Standard (IS) is included in the reaction mix, controlling the possible inhibition of the PCR reaction (ISIN version) and possibly also the DNA extraction process quality (ISEX version). IS positive amplification is detected in the HEX fluorophore fluorescence channel. The detection kit takes advantage of the "hot start" technology, minimizing non-specific reactions and assuring maximum sensitivity. Ready to Use MasterMix contains uracil-DNA-glycosylase (UDG), eliminating possible contamination of the PCR reaction by amplification products. The kit performs very sensitive L. pneumophila detection in clinical material (bronchoalveolar lavage, sputum). The kit is designed for in vitro diagnostics and provides qualitative detection.

### **ISIN** version

Internal Standard is included in the MasterMix tube. This PCR kit version enables PCR inhibition control.

### **ISEX** version

Internal Standard is provided as independent item within the package. This PCR kit version enables both PCR inhibition control and nucleic acid purificaion process efficiency control.

#### MICROBIOLOGICAL DNA DETECTION TECHNOLOGY ISIN ISEX / SAMPLE / SAMPLE positive sample contains Internal Standard is added into microorganism DNA ~~~ the sample before the DNA isolation / DNA ISOLATION / DNA ISOLATION the extracted sample DNA ~~ after the isolation the extracted is added into the Ready to use sample DNA ---- with MasterMix after the isolation the concurrently isolated Internal Standard DNA are added into and the tube is inserted into the real-time device the Ready to use MasterMix and the tube is inserted into the real-time device 4 + / PCR AMPLIFICATION / PCR AMPLIFICATION both pathogen DNA and Internal both pathogen DNA and Internal Standard DNA are amplified from Standard DNA are amplified from the same primer couple $\gtrsim$ the same primer couple $\gtrsim$ in the course of the PCR in the course of the PCR / EVALUATION

#### POSITIVE SAMPLE

- exponential fluorescence growth in the FAM fluorophore is evident if the detected pathogen DNA is present in the sample

#### QUALITY CONTROL FOR THE DIAGNOSTIC PROCESS

- exponential growth of the HEX fluorophore fluorescence, as a result of the IS amplification, controls the following:

- 1. Inhibition and efficiency of the PCR amplification -ISIN version
- 2. DNA extraction quality, inhibition and efficiency of the PCR amplification - ISEX version



# USER MANUAL

### SAMPLING AND SAMPLE STORAGE

*Legionella* detection in human clinical diagnostics is feasible from sputum or bronchoalveolar lavage samples. Sampling should be performed into sterile tubes without any transportation media and the samples should be transported to the laboratory within 12 hours at the temperature between +2 °C and +8 °C. In case of longer storage keep all samples frozen at the temperature between -85 and -10 °C.

### NUCLEIC ACID PURIFICATION

Nucleic acid isolation should be performed by isolation kits available at the market according to protocols for the particular clinical material isolation. The manufacturer recommends the following isolation kits:

GeneProof PathogenFree DNA Isolation Kit croBEE NA16 Nucleid Acid Extraction System

When using the ISEX versions of the PCR kits the IS should be added directly into the sample at the beginning of the isolation process so that in the end 1  $\mu$ l of the resulting elution volume contains 0.1  $\mu$ l of the IS:

Elution volume	25 µl	50 µl	100 µl	200 µl
Internal Standard	2,5 µl	5 µl	10 µl	20 µl

### PCR SETUP

1. Add 30 µl of MasterMix into PCR tubes.

2. Add 10  $\mu$ l of the isolated nucleic acid sample or 10  $\mu$ l of Positive Control into the individual PCR tubes. The final reaction mix volume will be 40  $\mu$ l.

It is necessary to keep all components at +2°C to +8°C during the PCR preparation.

3. Close the tubes, centrifuge shortly, insert them into the device and let them amplify according to the following PCR profile. Be very careful when handling the Positive Control or the clinical material, incorrect handling could result in contamination and the consequent impairment of the kit components or the MasterMix! The manufacturer is not responsible for the kit impairment due to incorrect handling.

### **AMPLIFICATION PROFILE**

Step	Temperature	Time	Data collection	Cycles
1. Hold	37 °C	2 min		1
2. Hold	95 °C	10 min		1
	95 °C	5 s		
3. PCR	60 °C	40 s	FAM+HEX	45
	72 °C	20 s		

# VALIDATED INSTRUMENTS

GeneProof PCR kits are designed for use with real-time devices from various manufacturers. This PCR kit has been validated with the following devices:

Applied Biosystems 7300/7500 Real-Time PCR System AriaMx Real-Time PCR System Dx/CFX96<sup>™</sup>/CFX Connect<sup>™</sup> Real-Time PCR Detection System LightCycler ® 2.0, LightCycler ® 480 LineGene 9600 Mx3000P/3005P QPCR System Rotor-Gene 3000, Rotor-Gene 6000, Rotor-Gene Q SLAN® Real-Time PCR System

GeneProof diagnostic kits are continually validated with various types of devices. Please request the current list at support@geneproof.com.



### **CLINICAL SAMPLES ANALYSIS EVALUATION**

Channel FAM	Channel HEX	Result	Interpretation	
	$\checkmark$	Valid	Legionella pneumophila	positive
		Valid	Legionella pneumophila	positive
	Ct<38	Valid	Legionella pneumophila	negative
	Ct>38	Invalid		
		Invalid		

### **CONTROL ANALYSIS EVALUATION**

	Channel FAM	Ch	annel HEX	Result
Positive Control		$\checkmark$	or	Valid
Negative Control		$\checkmark$	or	Valid
Positive Control		$\checkmark$	or	Invalid
Negative Control		$\checkmark$	or	Invalid

# WARNING

A single valid package insert for a specific kit is included in the package or to be requested for the particular lot from the manufacturer. The kit should be disposed of after use according to the current legal regulations considering the fact that the kit doesn't contain any dangerous, infectious or toxic components that would be subject to special safety regulations, and the packaging materials are made of paper and polypropylene. If you have any questions please contact our Customer Service.

Customer care and technical support

 Tel.:
 +420543211679

 Fax:
 +420516770824

 email:
 support@geneproof.com

Orders	
Tel.:	+420543211679
Fax:	+420516770824
email:	sales@geneproof.com



GeneProof a.s. Vídeňská 119 / CZ-619 00 Brno / +420 543 211 679 / info@geneproof.com Version: DOK\_142\_15\_01 Valid from: 20.07.2015





